

## Understanding Your Brain Tumor

The brain is a complex organ that controls all processes and functions of the body. Some of these functions include thought/memory, emotion, motor skills, temperature, vision, etc.

A brain tumor is a mass or growth of abnormal cells in your brain. These tumors can be cancerous (malignant) or noncancerous (benign). When you have a brain tumor, the part of the brain closest to the tumor may be damaged. This means that part of your brain is not able to function properly. The part of the brain that is involved or damaged will determine the symptoms you experience.

### Lobes of the brain and what they control:

#### Frontal:

- Personality
- Decision-making
- Movement
- Recognition of smell
- Speech/Language
- Emotions

#### Occipital

- Vision

#### Parietal

- Understanding where you are in space
- Ability to identify objects
- Interpreting pain and touch
- Understanding spoken Language

#### Temporal

- Short term memory
- Speech
- Musical rhythm
- Smell recognition

### Treatment of your Brain Tumor:

Treatment of your brain tumor can include radiation, chemotherapy, surgery, or a combination of these. Your surgeon has determined that including surgery in your treatment plan is the best option for you. The goal of surgery is to remove as much of the tumor as possible without affecting normal brain tissue and function. Removing the tumor can help resolve or lessen some of the symptoms you are experiencing.

### After Surgery

After surgery, you may need to stay in the hospital for a few days. When you are discharged following surgery, we expect that in most cases you will be self-sufficient in your personal care. However, depending on your home situation, arrangements may need to be made for additional help or potentially a transition to a short-term rehab center. This determination is made while you are in the hospital with the assistance of physical and occupational therapists. Additional arrangements for medical equipment for home use will also be made prior to leaving the hospital.

### Paperwork

Please do not bring any paperwork that needs to be completed by our office to the hospital with you. All paperwork related to disability, FMLA or other medical legal forms should be mailed to Midwest Spine and Brain Institute, Attention Medical Legal Department, 1950 Northwestern Avenue Suite 102. Stillwater, MN 55082. Please allow 1-2 weeks for forms to be completed. Midwest Spine & Brain Institute does charge an administrative fee for such paperwork to be completed. You may leave a message for our Medical Legal Department at 651-259-4522.

# Medication Instructions Before Surgery

You will need to stop taking certain medications and supplements before surgery.

If you have any questions, please contact our office at 651-430-3800.

## Blood Thinning Medication

Although it is not safe to proceed with surgery while taking the following “blood thinner” anticoagulant medication, it is also not safe to stop these medications without first consulting with the medical doctor who prescribed it for you. **If you are taking blood thinning medication, discuss a plan with the doctor who is prescribing the medication.**

These medications include, but are not limited to:

- Coumadin (warfarin)
- Plavix (clopidogrel)
- Eliquis (apixaban)
- Xarelto (rivaroxaban)
- Pradaxa (dabigatran)
- Brilinta (ticagrelor)

## Anti-Inflammatory Medication

It is recommended that you stop taking anti-inflammatory medications **10 days before surgery.** These medications include, but are not limited to the following:

Advil	Indocin
Aleve	Lodine
Anaprox	Motrin
Aspirin	Naproxen
Celebrex	Orudis
Daypro	Oruvail
Disalcid	Relafen
Ibuprofen	Vioxx
Excedrin	Meloxicam

## Herbal Supplements

It is recommended that you stop taking the following herbal supplements **7 days before surgery.**

Danshen	Ginseng
Dong quai	Goldenseal
Echinacea	Kava
Ephedra	Licorice
Feverfew	St. John’s Wort
Garlic	Valerian
Ginger	Vitamin E
Gingko	Yohimbe
Fish Oil	Any Multivitamin
Weight Loss Medications/Supplements	

## GLP-1 Agonists

It is recommended to hold the following daily or twice daily injections **one day prior to Surgery and day of surgery.**

- Exenatide (Byetta)
- Lixisenatide (Adlyxin)
- Liraglutide (Victoza, Saxenda)
- Lixisenatide with glargine (Soliqua)
- Wegovy
- Phentermine

For Weekly injections or daily oral dosing **Hold 7 days prior to surgery.**

- Albiglutide (Tanzeum)
- Exenatide (Bydureon)
- Dulaglutide (Trulicity)
- Semaglutide (Ozempic)
- Tirzepatide (GLP-1/GIPRA) (Mounjaro)
- Semaglutide (Rybelsus)

# Preoperative Instruction

**Dear Patient: Please take this preoperative instruction sheet to your family physician when you have your pre-op examination.**

Dear Health Care Provider,

Please perform pre-operative evaluation and testing listed below, which is based on patient's health condition(s) and current evidence-based guidelines. Thank you!

## **Lumbar Fusion (One or Two Levels), Lumbar Artificial Disc Replacement and Laminectomy (Cervical, Thoracic and Lumbar) to Include:**

- CBC
- POTASSIUM If on diuretics, anti-hypertensive or cardiac medications
- BMP If patient is taking an ACE/ARB medication
- EKG Male/Female 65 years of age or greater – needs within 1 year prior

## **Anterior/Posterior Spine Surgery (Cervical and Lumbar), Lumbar Spinal Fusion (3 or More Levels) and Thoracic Fusion to Include:**

- CBC
- INR
- BMP
- POTASSIUM If on diuretics, anti-hypertensive or cardiac medications
- EKG Male/Female 50 years of age or greater – needs within 1 year prior

## **Please see the following for any additional condition specific testing:**

- ❖ Cardiovascular Disease
  - CBC
  - BMP
  - EKG
  - CXR
- ❖ History of Stroke
  - EKG
- ❖ Bleeding Disorder/Anemia
  - CBC
  - INR
- ❖ Pulmonary Disease
  - CXR
- ❖ Hepatic Disease
  - CBC
  - INR
  - CMP
- ❖ Renal Disease
  - CBC
  - BMP
  - EKG
- ❖ Endocrine Disorder
  - BMP
  - EKG
- ❖ BMI >35 **AND** another risk factor
  - EKG
- ❖ Chemotherapy
  - CBC

1950 Northwestern Avenue Suite 102  
Stillwater, MN 55082-7615  
Phone: 651-430-3800  
Fax: 651-430-3827  
[www.midwestspineandbrain.com](http://www.midwestspineandbrain.com)

# Attention: Medical Facilities Performing Pre-Op Examinations

Please fax a copy of the pre-op exam to the hospital where the patient is having their surgery performed. Also, please fax a copy to Midwest Spine & Brain Institute. It is imperative that the hospital and Midwest Spine & Brain Institute have a copy before the surgery can be performed. Many times the surgery is early in the morning and the clinic where the pre-op was performed has not opened making it difficult to obtain a copy of the pre-op if necessary. Please ask the patient what particular hospital they will utilize.

**Please bring a copy of your pre-op physical to the hospital ONLY if your surgery is scheduled for the next day.**

**Midwest Spine & Brain Institute fax number: 651-430-3827**

## **Fax Numbers to commonly used Hospitals and Surgery Centers:**

<b>Fairview Ridges Hospital</b>	<b>952-892-2078</b>
<b>Fairview Ridges Surgery Center</b>	<b>952-898-3482</b>
<b>Greenway Surgery Center</b>	<b>612-728-2660</b>
<b>Maplewood Surgery Center</b>	<b>651-471-9748</b>
<b>North Memorial Health Hospital</b>	<b>763-581-3821</b>
<b>St. Croix Regional Medical Center</b>	<b>715-483-0519</b>
<b>St. John's Hospital (HealthEast)</b>	<b>651-326-8631</b>
<b>United Hospital</b>	<b>651-241-5073</b>
<b>Woodbury Surgical Suites</b>	<b>651-493-0344</b>

**Thank you for your assistance.**

# Recovery at Home after Surgery

## Recovery Time

The symptoms you had before surgery can take weeks or months to improve. It is common for these symptoms to come and go after surgery.

Be patient and allow your body time to heal.

## Bathing

For three to six weeks after surgery, or until your incision is healed, you can only take showers. Do not take baths.

- Cover your incision with plastic wrap and tape until drainage stops. Once your incision is no longer draining, you may shower with it uncovered.
- You may use a stool to sit on (like you did in the hospital).
- Keep your soap, shampoo and other items within reach.
- If you drop something, do not try to pick it up. Ask for help or have extra supplies handy.

## Dressings & Bandages

- If you have had Carpal Tunnel surgery, do not change your dressing until 2 days after surgery.
- If you have had Ulnar Nerve Transposition surgery, do not change your dressing until 3 days after surgery.
- If you have had spine surgery, change your dressing daily, using light gauze and tape, until drainage stops.
- Once drainage has stopped, you can leave your incision uncovered.
- Keep your incision clean and dry.
- Change your dressing after showering, and when it becomes wet.

## Stitches, Staples & Surgical Glue

- If you have staples along your incision, they will be removed at your follow-up visit two to three weeks after surgery.

- If you have small strips of tape (Steri-Strips) along your incision, they will fall off on their own within 2-3 weeks. Do not pull these off. When they fall off, they do not need to be replaced.
- If your wound was closed with dissolving sutures and your skin closed with surgical glue, you may get your skin wet immediately. You do not need to cover your incision.

## Call an Ambulance

**CALL 911 IF YOU ARE EXPERIENCING CHEST PAIN, SHORTNESS OF BREATH, OR DIFFICULTY BREATHING**

## Call Your Care Team

Call 651-430-3800 to speak with your care team if you have:

- a temperature of 101° F or higher
- yellow or green drainage or more than a slight amount of bloody drainage from your incision
- redness, swelling, or warmth by your incision
- new or unusual pain, numbness, or tingling
- pain you cannot control
- any bowel or bladder changes
- an opening in your incision
- pain in your calf or pressure in your legs
- been sent home from surgery with a drain
- any other questions or concerns

## Movement & Exercise

### Walking

- Initially, walk for exercise and to regain your strength and endurance. Begin slowly, and increase the amount you walk as tolerated.
- If you become sore or have pain, decrease the amount of walking for 1-2 days. Slowly increase the amount you walk again as tolerated.

### Sitting

- Practice good posture.
- Use fatigue as a reminder to rest.

- Increase the length of time you sit as necessary and as tolerated.

#### Bending, Twisting & Lifting

- Do not bend or twist a lot.
- If bending and lifting are necessary, do so at your knees and keep your back straight.
- Use a “reacher” to pick items up off the floor.
- Try not to lift items heavier than 5 pounds until your first follow up appointment.
- Carry items close to your body at waist level.
- Avoid activities like sweeping and vacuuming.

#### Steps

- Limit steps to only a few initially. You may gradually increase the number of steps you take as tolerated.
- Take your time. Steps may be challenging as surgery may make it hard to see where your feet are placed. Be sure your feet are firmly placed on each step before shifting your weight.

#### Driving

- Do not drive while taking prescription pain medication.

#### Sexual Activities

- You may resume sexual activities as your symptoms allow.

#### Bracing

- If you have had a hardware removal, spinal cord stimulator, carpal tunnel release, or ulnar nerve transposition, you typically will not need a brace. Your surgeon will let you know if one is needed.

#### Avoid Constipation

Prescription pain medicine, anesthesia and decreased activity can slow your digestion and cause constipation.

- Take stool softeners, such as Colace, as needed while you are taking narcotic pain medicine.
- Do not strain to have a bowel movement.
- You may try Milk of Magnesia or Miralax if stool softeners are not effective.
- Drink plenty of water.
- Include fruits and vegetables into your diet.

#### Prescription Pain Medication

Depending on your surgery and condition, your medicine will be refilled for a short time after surgery. If you still need pain medication after this time, your provider may refer you to your primary care physician or a pain clinic for further management.

- Take your recommended doses when your pain is at its worst. Slowly cut back (taper) on the narcotic when you think your pain is under control.
- Benzodiazepines (medications to calm or relax you) may need to be reduced or stopped before surgery and while taking any pain medication after surgery. Please discuss a plan for this with your primary care physician.
- Please call your care team if you have questions about your pain medication plan.

#### Questions

Call 651-430-3800 to speak with your care team.

# Potential Risks and Complications of Neurosurgery

Although the most likely complications are identified below, this is not a complete list, and other complications can occur. These risks should not be taken lightly; however, the possibility of any of these happening is very low.

- Your surgeon cannot guarantee relief of pain or other symptoms following surgery. Based on your surgical outcome, additional surgery/treatments may be necessary.
- X-ray may be used during your surgery. If you are pregnant, this could be unsafe for your baby.
- There is a risk for an allergic reaction to general anesthesia (being put to sleep).
- Lung problems (such as pneumonia), blood clots, heart attack, stroke, injury to a major blood vessel and wound infections can occur.
- Fluid or blood accumulation near the wound is possible. This can cause increased intracranial pressure, which can lead to visual problems, speech disturbance, memory problems, muscle weakness and/or loss of balance/coordination.
- Seizures can occur after surgery, regardless of if they were already present before surgery or not. Treatment is provided to help prevent seizures, though they still could occur with treatment.
- Cranial nerve palsies (lack of function of the cranial nerve) can occur. This may result in complete or partial weakness or paralysis of the areas of the body affected by a specific cranial nerve.
- If a spinal nerve is injured, it could result in permanent pain, numbness, or weakness in a limb, or loss of bowel or bladder control.
- If the spinal cord is injured, it could result in permanent paralysis in the legs and possibly parts of the arms.
- The spinal nerves and spinal cord travel through the vertebrae in a sac filled with spinal fluid. If the sac is punctured, spinal fluid will leak out. This is known as a dural leak. If this occurs you may be advised to remain flat in bed to prevent a spinal headache and allow the puncture site to heal. On rare occasions, an injection or further surgery may be required to seal the puncture site.
- A significant amount of blood can be lost during surgery, which may require you to have a blood transfusion. Your surgeon may recycle your own blood during surgery so, if needed, you may receive it back in a transfusion during or after surgery. We will help you make the necessary arrangements. If you do need blood from the blood bank, risks include hepatitis, allergic reactions, and on extremely rare occasions HIV/AIDS.

**I understand the potential benefits of the proposed treatment and any alternative forms of treatment, including my option of receiving no treatment. By signing below, I agree that I have had the opportunity to ask any and all questions that I may have and wish to proceed with surgery.**

**Patient (or Representative) Signature:** \_\_\_\_\_ **Date:** \_\_\_\_\_

**Witness Signature:** \_\_\_\_\_ **Date:** \_\_\_\_\_

## Notice of Disclosure

---

The following disclosures may apply to the treatment you receive at Midwest Spine & Brain Institute (MSBI). Please contact your MSBI physician or Administration if you have any questions about any of the following disclosures.

### **Disclosure of Ownership in Ambulatory Surgery Centers**

If your MSBI physician refers you to Maplewood Surgery Center, Woodbury Surgery Center or Greenway Surgery Center, the following disclosure applies:

Your health care provider is referring you to a facility or service in which your health care provider has a financial or economic interest. The MSBI physicians who are listed below have an ownership interest in these ambulatory surgery centers. You are free to choose a different ambulatory surgery center. If you would like to do so, please notify your physician and we will be happy to accommodate your request.

This disclosure applies to the following MSBI Physicians for Maplewood Surgery Center: David T. Chang, M.D., Ph.D.

This disclosure applies to the following MSBI Physicians for Greenway Surgery Center: Stefano M. Sinicropi, M.D.; Glenn R. Buttermann, M.D., M.S.; David T. Chang M.D. Ph.D.

This disclosure applies to the following MSBI Physicians for Woodbury Surgery Center: Glenn R. Buttermann, M.D., M.S.

This disclosure applies to the following MSBI Physicians for Valley Surgery Center: Eduardo Perez, M.D.

### **Disclosure of Ownership in Outside Facilities**

Your health care provider is referring you to a facility or service in which your health care provider has a financial or economic interest. The MSBI physician listed below has an ownership interest in this outside clinic. You are free to choose a different clinic. If you would like to do so, please notify your physician and we will be happy to accommodate your request.

This disclosure applies to the following MSBI Physicians for HyperCharge Clinic: Stefano M. Sinicropi, M.D.

### **Disclosure of Industry Relationships**

If you are scheduled to receive a neurosurgical or orthopedic implant, biologic product or device during your surgical procedure, it is possible that your spine surgeon may have a financial relationship with the manufacturer. Your surgeon will be glad to discuss the reasons for his/her choice of implant/product. You have the right to choose another facility or provider for your procedure. Glenn R. Buttermann, M.D., M.S., holds multiple patents. His industry consulting relationships include FG Solco. He also owns shares in Nexxt Spine LLC.

### **Insurance Coverage for Other Facilities**

Some facilities may be out of network for certain health plans. Please contact the facility as well as your insurance plan for details regarding your coverage.